

## IN VIVO AND IN VITRO ANTITUMOR EFFECT OF A UNIQUE NUTRIENT MIXTURE ON LUNG CANCER CELL LINE A-549

**M. Waheed Roomi, Vadim Ivanov, Tatiana Kalinovsky, Aleksandra Niedzwiecki, and Matthias Rath** □ *Dr. Rath Research Institute, Cancer Division, Santa Clara, California, USA*

□ *The high incidence of lung cancer and ineffective toxic action of current mono and doublet chemotherapy approaches result in poor patient survival. Further, matrix metalloproteinases (MMPs) are implicated in neoplastic invasion and metastasis. Based on this, the authors investigated the effect of a dietary micronutrient mixture (NM) containing lysine, proline, arginine, ascorbic acid, and green tea extract on the tumor growth of human lung carcinoma cell A-549 xenografts in athymic nude mice. Additionally, the authors tested the in vitro antitumor effect of NM on lung carcinoma A-549 cells by measuring cell proliferation by MTT assay, MMP-2 and -9 secretion by gelatinase zymography, and cell invasion through Matrigel. Nutrient supplementation strongly suppressed the growth of tumors without adverse effects in nude mice; tumor weight was reduced by 44% ( $P = .0001$ ) and tumor burden was reduced by 47% ( $P < .0001$ ) with supplementation. Zymography demonstrated in vitro secretion of MMP-2 by uninduced human lung carcinoma cells and both MMP-2 and -9 by phorbol 12-myristate 13-acetate (PMA) (200 ng/mL)-treated cells. NM inhibited the secretion of both MMPs in a dose-dependent fashion, with virtual total inhibition at 500 µg/mL concentration. The invasion of human lung carcinoma cells through Matrigel was significantly reduced at 100 µg/mL (64%) and totally inhibited at 500 µg/mL concentration of NM ( $P = .01$ ). Suppression of lung tumor growth in nude mice and inhibition of MMP secretion and Matrigel invasion suggest NM may act as an anticancer agent and as such warrants further investigation.*

**Keywords** A-549, antitumor effect, lung cancer, MMP, nutrient mixture

Lung cancer is the leading cause of cancer-related death for both men and women worldwide. For 2005, the American Cancer Society estimates approximately 172,570 new cases of lung cancer in the United States [1]. Worldwide incidence of lung cancer has increased from one million in 1999 to 1.35 million new cases in 2002 [2]. Despite a shift in treatment

Received 16 June 2006; accepted 9 August 2006.

Consulting pathologist Dr. Alexander De Paoli, DVM, PhD, provided the histology slides.

Address correspondence to Aleksandra Niedzwiecki, Dr. Rath Research Institute, Cancer Division, 1260 Memorex Drive, Santa Clara, CA 95050, USA. E-mail: a.nied@drath.com

strategy in the 1970s from radiotherapy to sequential chemoradiation to concurrent chemoradiation, only marginal improvement in survival has been seen; nearly 163,510 Americans will die from lung cancer this year and 5-year survival is limited to 5% to 10% [3]. Average age at diagnosis is 70 and the American Cancer Society estimates that nearly 6 out of 10 people with lung cancer will die within 1 year of diagnosis. Non-small-cell lung cancer (NSCLC), a heterogeneous aggregate of histologies, comprising squamous, adenocarcinoma, and large-cell carcinoma, is the dominant histology responsible for roughly 80% of all lung malignancies [4]. Approximately 50% of NSCLC cases present with incurable metastatic (stage IV) disease [5]. Not only does radiation therapy and adjunctive chemotherapy fail to improve survival in NSCLC patients, but these strategies are associated with dose-dependent toxicities due to nonselective damage to cancerous and normal cells. Cisplatin is associated with severe nephrotoxicity [6]; vincristine, with dose-limiting neurotoxicity, manifested mainly as peripheral neuropathy [7]; doxorubicin with cardiotoxicity, which can lead to acute and chronic heart failure [8, 9]; etoposide with bone marrow suppression and gastrointestinal toxicity [10]; and cyclophosphamide with significant pulmonary toxicity [11]. Taken together, late diagnosis of lung cancer, high mortality, and the ineffective and harmful effects of chemotherapy and radiotherapy necessitate the adoption of novel treatment approaches that target and arrest metastasis.

Proteolytic degradation of the extracellular matrix (ECM) is necessary for cancer cells to invade and metastasize. Matrix metalloproteinases (MMPs) are zinc-dependent endopeptidases implicated in facilitating tumor invasion and metastasis through perforation of physical barriers to invasion by degradation of ECM macromolecules and modulation of cell adhesion and activation of ECM components to expose hidden biologic activities. Due to this, recent research efforts have focused on the role of MMP inhibitors in the modulation of tumor invasion, metastasis, and angiogenesis. Rath and Pauling proposed that natural inhibitors, such as lysine and ascorbic acid, have the potential to inhibit tumor growth and expansion through the modulation of ECM proteolysis and optimization of connective tissue integrity [12]. Our earlier work based on this concept led to the development of a nutrient composition (NM) that has shown to be effective in achieving a complete inhibition of ECM invasion of several cancer cell lines *in vitro*, strong antimetastatic effects in animal models [13, 14], and demonstrated significant antiangiogenic and proapoptotic effects [15, 16].

In this report we investigated the antitumor potential of this NM-containing ascorbic acid, lysine, proline, arginine, and green tea extract on human lung carcinoma cells A-549 *in vivo* and *in vitro*.

## MATERIALS AND METHODS

### Cancer Cell Line and Culture

Human lung cancer cells A-549 obtained from ATCC (American Type Culture Collection, Rockville, MD) were maintained in Ham F12K medium, supplemented with 10% fetal bovine serum, 100 U/mL penicillin, and 100  $\mu$ g/mL streptomycin. The media and sera used were obtained from ATCC, and antibiotics (penicillin and streptomycin) were from Gibco BRL, Long Island, NY.

### Composition of Nutrient Mixture

Stock solution of the NM was composed of the following in the ratio indicated: vitamin C (as ascorbic acid and as Mg, Ca, and palmitate ascorbate) 700 mg; L-lysine 1000 mg; L-proline 750 mg; L-arginine 500 mg; *N*-acetylcysteine 200 mg; standardized green tea extract (80% polyphenol) 1000 mg; selenium 30  $\mu$ g; copper 2 mg; manganese 1 mg.

The NM was formulated based on targeting different physiological processes involved in cancer progression and metastasis at the cellular level. For example, the ECM integrity is dependent upon adequate collagen formation and its stability. In this aspect ascorbic acid and the amino acids lysine and proline are necessary for the formation and optimum structure of collagen fibers. Manganese and copper are also essential cofactors in collagen formation process. Collagen stability can be controlled by lysine [12] and also by *N*-acetylcysteine through its inhibitory effect on MMP-9 activity [17] and invasive activities of tumor cells [18]. Also, selenium has been shown to interfere with MMP expression and tumor invasion [19], as well as migration of endothelial cells through ECM [18]. Ascorbic acid, in addition to its critical role in ECM structure, has been shown to inhibit cancer cell division and growth through production of hydrogen peroxide, especially when applied in high doses [20, 21]. Green tea extract has shown to be a promising agent in controlling angiogenesis, metastasis, and other aspects of cancer progression [22]. Because arginine is a precursor of nitric oxide (NO), any deficiency of arginine can limit the production of NO, which has been shown to predominantly act as an inducer of apoptosis, as in the case of breast cancer cells [23].

Based on our own research and published data, we postulated that metabolic effects of a combination of ascorbic acid, lysine, proline, green tea extract, arginine, *N*-acetylcysteine, selenium, copper, and manganese would result from their synergy. For example, we found that a combination of ascorbic acid, lysine and proline used with epigallocatechin gallate (EGCG) enhanced the anti-invasive activity of 20  $\mu$ g/mL EGCG to that of

50 µg/mL [24]. Thus by including nutrients such as *N*-acetylcysteine, arginine, selenium, manganese, and copper in addition to ascorbic acid, proline, lysine, and EGCG, we could obtain significant reduction in cell invasion at a much lower concentration of EGCG.

## **In Vivo Studies**

### ***Animals***

Male athymic nude mice (NCr-nu/nu), approximately 6 weeks of age on arrival, were purchased from Simonsen Laboratories, Gilroy, CA, and maintained in microinsulator cages under pathogen-free conditions on a 12-hour light/12-hour dark schedule for a week. All animals were cared for in accordance with institutional guidelines for the care and use of experimental animals.

After housing for a week, the mice were inoculated subcutaneously with  $3 \times 10^6$  human lung cancer cells in 0.2 mL of phosphate-buffered saline (PBS) and 0.1 mL of Matrigel (BD Bioscience, Bedford, MA). After injection, the mice were randomly divided into 2 groups, A and B. Six mice were allocated to each group. From day 1, mice from group A were fed a regular diet and those in group B were fed a regular diet supplemented with 0.5% NM (*w/w*). During the study, the mice consumed, on the average, 4 g of their respective diets per day. (Thus, the supplemented mice received approximately 20 mg of NM per day.) After 4 weeks, mice were sacrificed; tumors were excised, weighed, and measured. Dimensions [length (L)  $\times$  width (W)] of the tumors were measured with a pair of digital calipers, and the tumor burden was calculated using the following formula:  $1/2 \times L \times W$ . Tumors were then fixed in 10% (*v/v*) buffered formalin and processed for histology.

### ***Histology***

Tissue samples were fixed in 10% buffered formalin. All tissues were embedded in paraffin and cut at 4 to 5 microns. Sections were deparaffinized through xylene and graduated alcohol series to water, and incubated for 5 minutes in aqueous 3% hydrogen peroxide to block endogenous peroxidase. Histological sections were stained with hematoxylin and eosin (H&E) for evaluation using a standard light microscope.

## **In Vitro Studies**

### ***Cell Culture***

At near confluence, the lung carcinoma A549 cells were treated with the NM, dissolved in medium and tested at 0, 10, 50, 100, 500, and 1000 µg/mL in triplicate at each dose. Cells were also treated with

200 ng/mL of phorbol 12-myristate 13-acetate (PMA). Five microliters of a 40  $\mu$ g/mL solution of PMA was added to 0.5 mL of medium. The plates were then returned to the incubator.

#### *MTT Assay*

Cell proliferation was evaluated by MTT [3-(4,5-dimethylthiazol-2-yl) 2,5-diphenyl tetrazolium bromide] assay 24 hours following incubation with test reagents. The MTT assay is a colorimetric assay based on the ability of viable cells to reduce a soluble yellow tetrazolium salt (MTT) to a blue formazan crystal by mitochondrial succinate dehydrogenase activity of viable cells. This test is a good index of mitochondrial activity and thus of cell viability. The cells were washed with PBS and 500  $\mu$ L of MTT (Sigma catalog number M-2128), 0.5  $\mu$ g/mL in medium, was added to each well. After MTT addition, the plates were covered and returned to the 37°C incubator for 2 hours, the optimal time for formazan product formation. Following incubation, the supernatant was carefully removed from the wells, the formazan product was dissolved in 1 mL DMSO, and absorbance was measured at 570 nm in Bio Spec 1601, Shimadzu spectrometer. The OD<sub>570</sub> of the DMSO solution in each well was considered to be proportional to the number of cells. The OD<sub>570</sub> of the control (treatment without supplement) was considered 100%.

#### *Gelatinase Zymography*

MMP secretion in conditioned medium was determined by gelatinase zymography. Gelatinase zymography was performed in 10% Novex precast sodium dodecyl sulfate (SDS)-polyacrylamide gel (Invitrogen Corporation) in the presence of 0.1% gelatin under nonreduced conditions. Culture medium (20  $\mu$ L) mixed with sample buffer was loaded and SDS-PAGE was performed with Tris glycine SDS buffer as described by the manufacturer (Novex). Samples were not boiled before electrophoresis. Following electrophoresis the gels were washed twice in 2.5% Triton X-100 for 30 minutes at room temperature to remove SDS. The gels were then incubated at 37°C overnight in substrate buffer containing 50 mM Tris-HCl and 10 mM CaCl<sub>2</sub> at pH 8.0, stained with 0.5% Coomassie Blue R250 in 50% methanol and 10% glacial acetic acid for 30 minutes, and destained. Protein standards were run concurrently and approximate molecular weights were determined by plotting the relative mobilities of known proteins.

#### *Matrigel Invasion Studies*

Invasion studies were conducted using Matrigel (Becton Dickinson) inserts in 24-well plates. Suspended in medium, human lung cancer A-549 cells were supplemented with nutrients, as specified in the design of the experiment and seeded on the insert in the well. Thus both the medium on the insert and in the well contained the same supplements.

The plates with the inserts were then incubated in a culture incubator equilibrated with 95% air and 5% CO<sub>2</sub> for 24 hours. After incubation, the medium from the wells were withdrawn. The cells on the upper surface of the inserts were gently scrubbed away with cotton swabs. The cells that had penetrated the Matrigel membrane and migrated onto the lower surface of the Matrigel were stained with Hematoxylin and Eosin and visually counted under the microscope.

### Statistical Analysis

The results were expressed as means  $\pm$  SD or SEM, as indicated in the results, for the groups. Data was analyzed by independent-sample *t* test.

## RESULTS

### In Vivo Studies


#### *Tumor Growth*

Nude mice supplemented for 4 weeks with NM in their diets, developed significantly smaller tumors than did the control nude mice. Tumors developed in NM supplemented mice demonstrated lower weight (by 44%,  $P = .0001$ , as shown in Figure 1A) and tumor burden (by 47%,  $P < .0001$ , as shown in Figure 1B) compared to those developed by the control group of mice. Nude mice from both groups showed no body weight loss over the study period. The initial mean weight of mice was  $20.6 \pm 0.7$  g; the mean post treatment weight of the control group of mice was  $23.2 \pm 1.02$  g and of the supplemented group was  $21.3 \pm 1.02$  g.

#### *Histology*

Sections from irregularly round, focally invasive subcutaneous masses of the control group of mice were composed of nests and sheets of irregularly round to cuboidal epithelioid cells with irregularly round nuclei and variably vacuolated cytoplasm consistent with a carcinoma. Bluish staining material, probably mucin, was occasionally seen in the interstitial tissue of tumors. Mitotic figures averaged about 1 per high-power field. Foci of tumor necrosis infiltrated with neutrophils were variably disseminated in the tumor, involving about 10% of the masses. Tumor masses were surrounded by fibroblasts and a moderate mixed inflammatory cell infiltrate. The histology of masses from the supplemented group were similar to the control group, except the necrosis involved about 20% of the tumor masses and the inflammatory cell response surrounding the tumors consisted primarily of macrophages.






**FIGURE 1** Effect of nutrient supplementation (NM 0.5% diet) on mean tumor weight (A) and mean tumor burden (B) in male nude mice (6 in each group) 4 weeks post inoculation with human lung cancer A-549 cell xenografts. Nutrient supplementation reduced tumor weight by 44% ( $P = .0001$ ) and mean tumor burden by 47% ( $P < .0001$ ). Error bars on graphs represent SEM. Tumor burden was calculated as  $1/2 \times L \times W$ .

## In Vitro Studies

### Cell Proliferation Study

The nutrient mixture applied at lower concentrations (up to 500  $\mu\text{g}/\text{ml}$ ) did not show any significant effect on lung carcinoma cell growth. Cell proliferation increased to 144% ( $P = .002$ ) over the control at NM 500  $\mu\text{g}/\text{mL}$ . However, at 1000  $\mu\text{g}/\text{mL}$ , NM inhibited growth to 80% of the control ( $P = .008$ ), as shown in Figure 2. It is difficult to explain the sudden decrease in cell proliferation from 500 to 1000  $\mu\text{g}/\text{mL}$ . It is possible that the cells were not washed properly and that the residual NM interfered with the assay. However, current studies in progress have demonstrated




**FIGURE 2** Effect of the nutrient mixture (NM) on cell proliferation of lung cancer A-549 cell line: MTT assay 24 h. NM showed minimal stimulation of A-549 cells up to 500 µg/mL, increased cell proliferation by 40% at 500 µg/mL, and reduction in cell growth to 80% of the control at 1000 µg/mL. Error bars represent SD.

increased apoptosis of A549 cells with higher NM concentration, especially at 500 and 1000 µg/mL.


#### *Gelatinase Zymography Study*

Zymography assays have shown that uninduced human lung carcinoma cells secrete only MMP-2 (Figure 3A), whereas PMA treatment (200 ng/mL) results in secretion of both MMP-2 and -9 (Figure 3B). Exposure of cells to NM resulted in inhibition of the secretion of both MMPs in a dose-dependent fashion, with virtual total inhibition at 500 µg/mL concentration.



**FIGURE 3** Effect of the nutrient mixture (NM) on lung carcinoma A-549 MMP-2 and MMP-9 secretion (A) and on PMA (200 ng/mL)-treated lung carcinoma A-549 MMP-2 and MMP-9 secretion (B). Lane 1, markers; lane 2, control; lanes 3–7, NM 10, 50, 100, 500, 1000 µg/mL. Zymography demonstrated secretion of MMP-2 and PMA-induced MMP-9 secretion. NM inhibited the secretion of both MMPs in a dose-dependent fashion with total inhibition at 500 µg/mL.





**FIGURE 4** Effect of the nutrient mixture (NM) on lung cancer A-549 Matrigel invasion. NM inhibited Matrigel invasion of A-549 cells by 64% at 100 µg/mL and totally blocked it at 500 µg/mL NM ( $P < .001$ ). Error bars represent SD.

### *Invasion Study*

The invasion of human lung carcinoma cells through Matrigel was significantly reduced at 100 µg/mL (64%) and totally inhibited at 500 µg/mL concentration of the nutrient mixture ( $P = .01$ ), as shown in Figure 4.

## **DISCUSSION**

Matrix metalloproteinases are involved in tumor growth, angiogenesis, invasion, and metastasis. Tumor growth depends upon the generation of new blood vessels to sustain survival and proliferation. Growing tumors become hypoxic and acidotic beyond the size of 2 mm and secrete several growth factors to stimulate local blood vessels to sprout branches. Activation of endothelial cells give rise to the production of matrix metalloproteinases, allowing for the decomposition of the surrounding extracellular matrix and eventual development into new blood vessels.

The results of this *in vivo* study of human lung cancer cell xenografts in immune impaired (athymic) male nude mice demonstrated significant suppression of lung tumor growth (44% reduction in tumor weight and 47% reduction in tumor burden) with nutrient supplementation (NM 0.5%). In a previous study we showed that NM caused a significant ( $P < .05$ ) reduction in basic fibroblast growth factor (bFGF)-induced angiogenesis in a chorioallantoic membrane (CAM) assay in chick embryos at a

concentration of 250  $\mu\text{g}/\text{mL}$ , as well as decreased human osteosarcoma U2OS cell expression of Vascular endothelial growth factor (VEGF), angiopoietin-2, bFGF, Platelet-derived growth factor (PDGF), and transforming growth factor (TGF)- $\beta$ -1 [15]. Though the results from the cellular proliferation study were questionable due to the large difference between the effect at 500 and 1000  $\mu\text{g}/\text{mL}$ , it is only one of the parameters involved in tumor growth, as indicated above. Furthermore, studies in progress indicate that NM causes increased apoptosis in lung cancer A549 cells with higher doses, especially at NM 500 and 1000  $\mu\text{g}/\text{mL}$ . In addition, a study on the effects of this nutrient mixture on human T-lymphocytic virus (HTLV)-1-positive (HuT-102 and C9-PL) and-negative (CEM and Jurkat) cells revealed that this combination of nutrients significantly increased apoptotic cells in the preG<sub>1</sub> phase, secondary to up-regulation of p53, p21, and Bax protein levels and down-regulation of Bcl-2 $\alpha$  protein expression [16].

Control of proteolytic activity of ECM provides an opportunity of addressing common mechanism of metastasis, angiogenesis and tumor growth. Research has shown that highly metastatic lung cancer and other cancer cells secrete higher amounts of MMPs than do poorly metastatic cells. Therefore, prevention of ECM degradation through the inhibition of MMP activity, specifically MMP-2 (gelatinase A) and MMP-9 (gelatinase B) has been shown to be a promising therapeutic target in blocking tumor progression. Rath and Pauling postulated [12] that cancer growth and metastasis can be controlled through a natural approach, targeting the stability of the connective tissue surrounding cancer and therefore contributing to tumor "encapsulating" effect, using nutritional components, such as lysine and lysine analogues. Lysine interferes with the activation of plasminogen into plasmin by tissue plasminogen activator (tPA) by binding to plasminogen active sites, and consequently affecting the plasmin-induced MMP activation cascade [12]. A recent study demonstrated 7-fold reduction in metastasis of mammary cancer in plasmin deficient transgenic mice [25]. Lysine-mediated effects on the ECM include increased connective tissue strength and stability. It is well known that optimization of synthesis and structure of collagen fibrils depends upon hydroxylation of proline and lysine residues in collagen fibers catalyzed by ascorbic acid. Suboptimal levels of ascorbic acid and lysine are possible in various pathological stages and in deficient diets as these nutrients are not produced in the human body.

As mentioned previously, MMPs, especially MMP-2 and MMP-9, are also involved in tumor metastasis. Currently there is no effective treatment to blocking lung metastasis, and discovery of antimetastatic agents, especially nontoxic ones, would dramatically improve lung cancer prognosis. Invasion of host tissues is a hallmark of metastasis, a process dependent on tumor

cell adhesion, cell migration, and proteolytic degradation of the ECM by MMPs [26]. The results from our *in vitro* study of human lung cancer cells demonstrated potent, significant suppression of MMP secretion and Matrigel invasion and migration of A-549 cells at 500  $\mu\text{g}/\text{mL}$ . These results corroborate with our earlier data obtained with various types of cancer cell lines, indicating that NM affects mechanisms involved in cellular invasion that are universal for various types of cancers rather than being cancer specific [13–16].

The inhibitory effects of the individual nutrients tested have been reported in both clinical and experimental studies. Ascorbic acid has been reported to exert cytotoxic and antimetastatic actions on malignant cell lines [21, 27–30]; in addition, low levels of ascorbic acid have been reported in cancer patients [31, 32]. Green tea extract is a potent anticancer agent that has been reported to have antimetastatic and anti-angiogenic effects against human cancer cell lines; however, its effects were observed at much higher doses than available in food sources [33–35]. In this aspect, our previous studies indicated that the inhibitory effect of ascorbic acid, proline, lysine, and EGCG on several cancer cell lines in tissue culture studies was greater than that of the individual nutrients, allowing for obtaining maximum effects using lower doses of individual components [24].

Over the years, cancer therapy has witnessed many exciting developments, but a cure for cancer still remains to be found, because therapeutic approaches have relied mainly on killing tumors, without fully realizing the mechanisms controlling cancer cell death. Shifting research directions towards controlling cancer cell invasiveness in the tissues and their growth opens up a possibility of finding effective controls of this disease. Application of natural components in mixtures, not individually, can assure enhanced effectiveness on multiple targets.

Furthermore, in contrast to the toxic side effects of current mono and doublet chemotherapy, the nutrient mixture has been shown to be safe therapeutic agent. In a previous *in vivo* study addressing safety issues, we found that gavaging adult female ODS rats (weighing 250 to 300 g) with the nutrient mixture (at 30, 90, or 150 mg per day for 7 days), had neither adverse effects on vital organs (heart, liver, and kidney), nor on the associated functional serum enzymes, indicating that this mixture is safe to use even at these high doses, which far exceed the normal equivalent dosage of the nutrient [36]. Specifically, no difference was observed in liver serum levels of protein, albumin (A), globulin (G), A/G ratio, alkaline phosphatase, and aspartate aminotransferase (AST) between treated and control groups. However, alanine aminotransferase slowly decreased with increased dose of NM. In regard to heart serum levels, lactate dehydrogenase, creatine kinase, and AST in test groups did not differ from control groups. In regard to renal function, creatinine content (C) was the same for the test

and control groups, and blood urea nitrogen (BUN) (B) showed a decrease, as did the B/C ratio, with an increased concentration of NM. Furthermore, in regard to lipid levels, in the NM groups, both cholesterol and triglyceride levels decreased compared to control groups. There was no difference in the histopathology of liver, kidney, heart, and lung compared to control groups and amongst test groups.

## REFERENCES

- [1] American Cancer Society: Cancer Facts and Figures 2005. Atlanta, GA: American Cancer Society.
- [2] Parkin DM, Bray F, Ferlay J, Pisani P: Global cancer statistics, 2002. *CA Cancer J Clin.* 2005;55:74-108.
- [3] Lassen U, Osterlind K, Hansen M, Dombernowsky P, Bergman B, Hansen HH: Long-term survival in small-cell lung cancer: posttreatment characteristics in patients surviving 5 to 18+ years—an analysis of 1,714 consecutive patients. *J Clin Oncol.* 1995;13:1215-1220.
- [4] Potosky AL, Saxman S, Wallace RB, Lynch CF: Population variations in the initial treatment of non-small-cell lung cancer. *J Clin Oncol.* 2004;22:3261-3268.
- [5] Souquet PJ, Tan EH, Rodrigues Pereira J, Van Klaveren R, Price A, Gatzemeier U, Jaworski M, Burillon JP, Aubert D: GLOB-1: a prospective randomised clinical phase III trial comparing vinorelbine-cisplatin with vinorelbine-ifosfamide-cisplatin in metastatic non-small-cell lung cancer patients. *Ann Oncol.* 2002;13:1853-1861.
- [6] Xiao T, Choudhary S, Zhang W, Ansari NH, Salahudeen A: Possible involvement of oxidative stress in cisplatin-induced apoptosis in LLC-PK1 cells. *J Toxicol Environ Health.* 2003;66:469-479.
- [7] Quasthoff S, Hartung HP: Chemotherapy-induced peripheral neuropathy. *J Neurol.* 2002;249:9-17.
- [8] Doyle JJ, Neugut AI, Jacobson JS, Grann VR, Hershman DL: Chemotherapy and cardiotoxicity in older breast cancer patients: a population-based study. *J Clin Oncol.* 2005;23:8597-8605.
- [9] Guerra J, De Jesus A, Santiago-Borrero P, Roman-Franco A, Rodriguez E, Crespo MJ: Plasma nitric oxide levels used as an indicator of doxorubicin-induced cardiotoxicity in rats. *Hematol J.* 2005;5:584-588.
- [10] Yanez JA, Teng XW, Roupe KA, Fariss MW, Davies NM: Chemotherapy induced gastrointestinal toxicity in rats: involvement of mitochondrial DNA, gastrointestinal permeability and cyclooxygenase-2. *J Pharm Sci.* 2003;6:308-314.
- [11] Kachel DL, Martin WJ 2nd: Cyclophosphamide-induced lung toxicity: mechanism of endothelial cell injury. *J Pharmacol Exp Ther.* 1994;268:42-46.
- [12] Rath M, Pauling L: Plasmin-induced proteolysis and the role of apoprotein(a), lysine and synthetic analogs. *Orthomol Med.* 1992;7:17-23.
- [13] Roomi MW, Roomi N, Ivanov V, Netke S, Kalinovsky T, Niedzwiecki A, Rath M: In vivo and in vitro antitumor effect of ascorbic acid, lysine, proline and green tea extract on human melanoma cell line A2058. *In Vivo* 2006; 20:25-32.
- [14] Roomi MW, Roomi NW, Ivanov V, Kalinovsky T, Niedzwiecki A, Rath M: Modulation of *N*-methyl-*N*-nitrosourea induced mammary tumors in Sprague-Dawley rats by combination of lysine, proline, arginine, ascorbic acid and green tea extract. *Breast Cancer Res.* 2005;7:R291-R295.
- [15] Roomi MW, Roomi N, Ivanov V, Kalinovsky T, Niedzwiecki A, Rath M: Inhibitory effect of a mixture containing ascorbic acid, lysine, proline and green tea extract on critical parameters in angiogenesis. *Oncol Rep.* 2005;14:807-815.
- [16] Harakeh S, Diab-Assaf M, Niedzwiecki A, Khalife J, Abu-El-Ardat K, Rath M: Apoptosis induction by Epican Forte in HTLV-1 positive and negative malignant T-cells. *Leuk Res.* 2006;30:869-881.
- [17] Kawakami S, Kageyama Y, Fujii Y, Kihara K, Oshima H: Inhibitory effects of *N*-acetyl cysteine on invasion and MMP 9 production of T24 human bladder cancer cells. *Anticancer Res.* 2001;21:213-219.

- [18] Morini M, Cai T, Aluigi MG, Noonan DM, Masiello L, De Floro S, D'Agostinin F, Albini A, Fassina G: The role of the thiol *N*-acetyl cysteine in the prevention of tumor invasion and angiogenesis. *Int J Biol Markers* 1999;14:268-271.
- [19] Yoon SO, Kim MM, Chung AS: Inhibitory effects of selenite on invasion of HT 1080 tumor cells. *J Biol Chem*. 2001;276:20085-20092.
- [20] Maramag C, Menon M, Balaji KC, Reddy PG, Laxmanan S: Effect of vitamin C on prostate cancer cells in vitro: effect on cell number, viability and DNA synthesis. *Prostate*. 1997;32:188-195.
- [21] Chen Q, Espey MG, Krishna M, Mitchell JB, Corpe CP, Buettner GR, Shacter E, Levine M. Pharmacologic ascorbic acid concentrations selectively kill cancer cells: action as a pro-drug to deliver hydrogen peroxide to tissues. *Proc Natl Acad Sci USA*. 2005;102:13604-13609.
- [22] Hare Y: *Green Tea: Health Benefits and Applications*. New York, Basel: Marcel Dekker; 2001.
- [23] Cooke JP, Dzau VJ: Nitric oxide synthase: role in the genesis of vascular disease. *Annu Rev Med*. 1997;48:489-509.
- [24] Roomi MW, Ivanov V, Niedzwiecki A, Rath M: Synergistic antitumor effect of ascorbic acid, lysine, proline, and epigallocatechin gallate on human fibrosarcoma cells HT-1080. *Ann Cancer Res Ther*. 2004;12:148-157.
- [25] Almholt K, Lund LR, Rygaard J, Nielsen BS, Dano K, Romer J, Johnson M: Reduced metastasis of transgenic mammary cancer in urokinase-deficient mice. *Int J Cancer*. 2005;113:525-532.
- [26] Duffy MJ: The role of proteolytic enzymes in cancer invasion and metastasis. *Clin Exp Metastasis* 1992;10:145-155.
- [27] Naidu KA, Karl RC, Coppola D: Antiproliferative and proapoptotic effect of ascorbyl stearate in human pancreatic cancer cells: association with decreased expression of insulin-like growth factor 1 receptor. *Dig Dis Sci*. 2003;48:230-237.
- [28] Anthony HM, Schorah CJ: Severe hypovitaminosis C in lung-cancer patients: the utilization of vitamin C in surgical repair and lymphocyte related host resistance. *Br J Cancer*. 1982;46:354-367.
- [29] Koh WS, Lee SJ, Lee H, Park C, Park MH, Kim WS, Yoon SS, Park K, Hong SI, Chung MH, Park CH: Differential effects and transport kinetics of ascorbate derivatives in leukemic cell lines. *Anticancer Res*. 1998;18:2487-2493.
- [30] Glaab WE, Hill RB, Skopek TR: Suppression of spontaneous and hydrogen peroxide-induced mutagenesis by the antioxidant ascorbate in mismatch repair-deficient human colon cancer cells. *Carcinogenesis*. 2001;22:1709-1713.
- [31] Nunez C, Ortiz de Apodaca Y, Ruiz A: Ascorbic acid in the plasma and blood cells of women with breast cancer. The effect of consumption of food with an elevated content of this vitamin. *Nutr Hosp*. 1995;10:368-372.
- [32] Kurbacher CM, Wagner U, Kolster B, Andreotti PE, Krebs D, Bruckner HW: Ascorbic acid (vitamin C) improves the antineoplastic activity of doxorubicin, cisplatin and paclitaxel in human breast carcinoma cells in vitro. *Cancer Lett*. 1996;103:183-189.
- [33] Cao Y, Cao R: Angiogenesis inhibited by drinking tea. *Nature*. 1999;398:381.
- [34] Ahmad N, Feyes DK, Nieminen AL, Agarwal R, Mukhtar H: Green tea constituent epigallocatechin-3-gallate and induction of apoptosis and cell cycle arrest in human carcinoma cells. *J Natl Cancer Inst*. 1997;89:1881-1886.
- [35] Garbisa S, Sartor L, Biggin S, Salvato B, Benelli R, Albini A: Tumor gelatinases and invasion inhibited by the green tea flavanol epigallocatechin-3-gallate. *Cancer*. 2001;91:822-832.
- [36] Roomi MW, Ivanov V, Netke SP, Niedzwiecki A, Rath M: Serum markers of the liver, heart, and kidney and lipid profile and histopathology in ODS rats treated with nutrient synergy. *J Am Coll Nutr*. 2003;22:477, Abstract 86.