



M Waheed Roomi, Bilwa Bhanap,
Aleksandra Niedzwiecki* and
Matthias Rath

Oncology Division, Dr. Rath Research Institute, Santa Clara, CA-95050, USA

Received: 02 February, 2019

Accepted: 08 April, 2019

Published: 09 April, 2019

*Corresponding author: Aleksandra Niedzwiecki, Ph.D., Oncology Division, Dr. Rath Research Institute, 1260 Memorex Drive, Santa Clara, CA-95050, USA, Tel: 408-567-5050; E-mail: author@drath.com

Keywords: Mesothelioma; MSTO-211H; Apoptosis; Cytotoxicity; MTT; LIVE green caspases

<https://www.peertechz.com>



Research Article

A Novel Nutrient Mixture Induces Apoptosis in Human Mesothelioma Cells (MSTO-211H) via Activation of Caspases

Abstract

Background: Malignant mesothelioma is a highly aggressive and fatal cancer of older people. Consistently associated with asbestos exposure, mesothelioma is diagnosed when it is extensively metastasized and it has a dismal prognosis.

Purpose of the study: Surgery, chemotherapy, and radiotherapy are the mainstay of treatment yet they are ineffective in increasing the mesothelioma patient survival. Therefore, a different approach is needed. A novel nutrient mixture containing green tea extract, ascorbic acid, lysine, and proline exhibited anti-cancer effects in various cancers. In our previous studies, the nutrient mixture was seen to block MMP secretion and invasion through Matrigel™ by mesothelioma cells. In the current study, we wanted to explore if the nutrient mixture could induce apoptosis in mesothelioma cells.

Results and main findings: The mesothelioma MSTO-211H cells were cultured in complete Ham F-12K medium and the cells were then treated with the nutrient mixture at 0-1000 µg/ml concentration. Cell cytotoxicity was measured by MTT assay, morphology by H&E staining, and the apoptosis by LIVE Green Caspases. The nutrient mixture did not show significant inhibition of cell proliferation. However, H&E staining at the dose of 100 µg/ml showed a few cellular changes, while significant changes pertaining to apoptosis morphology were observed at 500 and 1000 µg/ml. LIVE Green Caspases analysis showed cells in early and late apoptosis with increasing doses of the nutrient mixture.

Conclusions: Our results suggest that the nutrient mixture may provide a new supplemental strategy for enhancing the therapeutic options for mesothelioma

Abbreviations

NM: Nutrient Mixture; MMP: Matrix Metalloproteinases; FBS: Fetal Bovine

Background

Malignant mesothelioma is an aggressive tumor commonly originating in the pleura and less commonly in the peritoneum. It is more frequently diagnosed in men over 70 and seen less frequently in women. Approximately 3000 people are diagnosed with mesothelioma every year in the United States [1]. Although overall incidence of mesothelioma has been decreasing in the USA over the past decade, it is still a major cause of death in other parts of the world. Exposure to asbestos is one of the main risk factors for the development of mesothelioma. Occupations in the mining, textile, construction and ship building industries are some of the professions at high risk for asbestos exposure. Some reports suggest that treatments such as radiotherapy can also lead to mesothelioma [2]. Patients with mesothelioma generally do not have complete resolution

after surgery. Chemotherapy or radiotherapy treatments possess their own challenges and associated side effects. Newer treatment options such as photodynamic and gene therapy, and interleukin and interferon treatment are advocated in some cases of mesothelioma [1]. However the response is limited. There is an urgent need for a safe and alternative option to control this malignancy.

Due to their antitumor actions including induction of apoptosis, a number of plant-based phytochemicals are increasingly being used as important treatment cancer treatment methods. Our aim was to address cancer metastasis by using a unique mixture of nutrients using green tea extract, ascorbic acid, lysine, and proline. This nutrient mixture (NM) has demonstrated anticancer properties in a number of cancer cell lines both in vivo and in vitro. Our previous studies have shown that the NM can inhibit the expression of matrix metalloproteinases (MMP) enzymes and inhibit cell invasion of the Matrigel™ [3]. In the present study we evaluated if the antitumor effects of NM are because of the apoptosis induction.

Materials and Methods

Chemicals and reagents

MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide) Assay, fetal bovine serum (FBS), and streptomycin and penicillin antibiotics were obtained from Sigma Aldrich (St. Louis, MO). The LIVE green caspase kit was purchased from Molecular Probe Inc. (Eugene, OR). The Ham F-12K medium was obtained from Gibco (Long Island, NY). Cell culture plates were obtained from Co-star (Cambridge, MA). All other reagents were high grade and purchased from Sigma Aldrich.

Cell Culture and composition of the Nutrient Mixture (NM)

Human mesothelioma cells MSTO-211H were purchased from ATCC (American Type Culture Collection) (Rockville, MD). The cells were cultured in Ham-F12K media supplemented with 10% FBS and antibiotics (100 U/mL penicillin and 100 µg/ml streptomycin) at 37° C in 5% carbon dioxide, 95% oxygen in a humidified cell incubator. The media was changed every alternate day. Cells were plated in 24-well tissue culture plates at the density of 3×10^4 cells until confluent. Then the cells were treated with NM at concentrations of 0, 50, 100, 500, and 1000 µg/ml in triplicate. The composition of NM is indicated in table 1.

MTT cytotoxic assay

Cell viability was determined using MTT assay. The assay is a colorimetric assay based on the ability of the viable cells to reduce soluble yellow tetrazolium salts to blue formazine crystals. After incubation of the cells with NM for 24 hours, 50 µL of MTT reagent in media (0.5 mg/ml) was added to each well, and further incubated for 2 hours at 37° C. The solution was carefully aspirated and the blue formazine precipitate was dissolved in 1 ml DMSO and the absorbance (OD) was measured on a microplate reader at a wavelength of 570 nm in a BioSpec 1601 Shimadzu spectrometer. The percentage of inhibition was calculated.

H&E staining

The MSTO-211H cells were cultured in 24-well plates and were treated with NM in test concentrations at 0, 50, 100, 500 and 1000 µg/ml, respectively. After 24-hour incubation, the cells were washed with phosphate-buffered saline (PBS), fixed with methanol, and then stained with hematoxylin and eosin (H&E), and images were captured by microscope.

Apoptosis and LIVE green caspase assay

Caspase activity was evaluated by LIVE Green Caspase Kit. The cells were treated with NM at 0, 50, 100, 500, and 1000 µg/ml for 24 hours, near confluence. After washing with PBS, the cells were treated with caspase reagent as specified in the manufacturer's instructions (Molecular Probes Image-IT LIVE Green Poly Caspases Detection Kit 135104, Invitrogen). The MSTO-211H cells were then photographed under the fluorescent

microscope and counted. Green colored cells represent viable cells, orange and red colors represent early and late apoptotic cells, respectively.

Statistical analysis

The results were expressed as mean \pm standard deviation (SD) for the groups. Data was analyzed by the independent t-test.

Results

Cytotoxicity

The cells were treated with NM at various concentrations and the cell viability was assessed by MTT assay. Cytotoxic effect of NM and viability of the MSTO-211H mesothelioma cells were studied and the results are shown in figure 1. The NM had no effect at 50 µg/ml, however, the cell proliferation was inhibited by 10%, 15%, and 20% at NM concentrations 100, 500, and 1000 µg/ml, respectively.

H&E staining and morphology

Apoptotic pattern was studied after H&E staining of the mesothelioma MSTO-211H cells at different concentrations of the NM. The characteristic apoptotic morphology was observed indicated by features such as shrinkage of the cytoplasm, and darkly stained nuclei with intensely acidophilic cytoplasm. These changes were seen to be increasing in a dose-dependent fashion, that is, minor apoptotic changes were noticed at 100 µg/ml, and moderate to significant changes were seen at 500 and 1000 µg/ml of NM. (Figures 2A through 2E)

Table 1: Nutrient Composition of Nutrient Mixture (NM).

Components of NM	Weight (mg)
Vitamin C (as ascorbic acid, Mg and Ca ascorbate and ascorbyl palmitate)	700 µM
L-Lysine	1000
L-Proline	750
L-Arginine	500
N-Acetyl cysteine	200
Standardized green tea extract EGCG (epigallocatechin gallate) (80% polyphenol)	1000
Selenium	30
Copper	2
Manganese	1
Quercetin dihydrate	50




Figure 1: Effect of NM on cell viability: MTT assay using mesothelioma MSTO-211H cells treated with 0, 50, 100, 500, and 1000 µg/ml concentrations of NM respectively.

Apoptosis

Figures 3A through 3E show the analysis of apoptosis by LIVE Green Caspase using mesothelioma MSTO-211H cells treated with NM at 0, 100, 500, and 1000 µg/ml. Very few cells were seen in the early stages of apoptosis at the NM dose of 50 µg/ml. However, as the concentration of the NM increased to 500 and 1000 µg/ml, the cells seen in early as well as late stages of apoptosis also increased. Quantitative analysis of the data revealed the percentage of apoptotic cells increasing with an increased dose of NM as shown in table 2. Forty percent of the cells were in late stages of apoptosis at 100 µg/ml of NM, 60% were in late apoptosis at 500 µg/ml, and 84% of the MSTO-211H cells were seen in late stages of apoptosis at 1000 µg/ml of NM (Figure 3F).

Discussion

Although the pathogenesis of highly aggressive malignant mesothelioma remains unclear, it is widely considered to be a fatal cancer with an average survival of 4 to 18 months after diagnosis [1]. This is mainly due to the aggressiveness of cancer and the lack of effective treatment options for mesothelioma. In our previous study, the NM demonstrated dose-dependent decrease in MMP secretion and dose-dependent increase in TIMP-2 secretion by mesothelioma cell lines [4].

The development of any abnormal cell to grow into a cancerous tumor is dependent on several factors, one of which




Figure 3: Effect of NM on apoptosis of mesothelioma MSTO-211H cells. 3A-3E - Photomicrographs of cells treated with 0, 50, 100, 500, and 1000 µg/ml concentrations of NM respectively.




Figure 2: 2A through 2E- Effect of NM on morphological changes: H&E staining of mesothelioma MSTO-211H cells treated with 0, 50, 100, 500, and 1000 µg/ml concentrations of NM respectively.




Figure 3f: 3F- Analytical representation of the differential distribution of MSTO-211H cells in early or late phases of apoptosis upon treatment with 0, 50, 100, 500 and 1000 µg/ml NM concentrations.

Table 2: Quantitative analysis of the data showing percentage of apoptotic MSTO-211H cells compared to the dose of NM.

NM dose	Live cells	Early Apoptosis	Late Apoptosis
0 µg/ml	100%	2%	0%
50 µg/ml	82	5%	13%
100 µg/ml	40%	22%	38%
500 µg/ml	18%	22%	60%
1000 µg/ml	8%	8%	84%

is its ability to escape apoptosis. Apoptosis, also known as programmed cell death, is a complex process that occurs in several pathological situations. Cancer cells not only multiply uncontrollably, they successfully dodge this natural mechanism

of death. Various methods have been developed to study apoptosis using up-regulation and down-regulation of specific genes such as the Bax and p53 genes [5]. Cancer cells hijack the cell machinery and stop apoptosis by blocking the activity of a particular gene, p53. This protector gene first stops the division of abnormal cells, and then initiates either their repair or their destruction if the damage is severe. Normally, the p53 gene activates specific apoptotic proteins from the Bcl2 family and the caspases enzymes that initiate and continue the process of apoptosis. Caspase enzyme activation is one of the earlier stages of apoptosis. Pro-apoptotic signals initiate specific reactions within the cell that activate the caspase enzymes and result in cleavage of protein substrates followed by the disassembly of the cell. The stimulation of suppressed apoptotic pathways in cancer cells and the induction of apoptosis is a predominant mechanistic approach to target cancer. The study of apoptosis by activation of caspase enzymes is an emerging area of research. The family of caspase aspartate, specifically cysteine proteases, is emerging as playing a central role in apoptosis. Some examples of these important caspases are caspase -3, -7, -8, -9, and -10 [6,7]. Several drugs, including Aspirin and other nonsteroidal anti-inflammatory drugs (NSAIDs), are being researched in cancer cells for their usage as apoptosis inducers.

In this study we evaluated the effect of NM on caspase activity in mesothelioma cells MSTO-211H. We observed a consistent increase in caspase activity with increasing doses of NM from 100 µg/ml, and 500 µg/ml to 1000 µg/ml. The dose-dependent increase in caspase activity progressed and was evident in the form of cell apoptosis as seen with H&E staining as well. Moderate apoptosis was noticed at 100 µg/ml, which increased to significant levels at 1000 µg/ml with NM showing 82% apoptosis.

Based on our previous studies, we have developed a multi-targeted approach to address cancer growth, and its spread. We have seen that the NM is effective in inhibiting invasion and metastasis, and inducing apoptosis by using an innovative combination of synergistic nutrients that include natural substances like ascorbic acid, lysine, proline, and green tea extract. The NM used in the study was specifically developed to combine the individual anti-tumorigenic and pro-apoptotic properties of the component micronutrients. Either used alone, or in combination with lysine and proline, ascorbic acid has shown anti-cancer actions due to its critical role in collagen synthesis, and formation of the basement membrane of extra cellular matrix [9-15]. Production of strong collagen fibers and structurally strong basement membrane are essential in preventing the tumor growth and micro-metastases. Epigallocatechin gallate (EGCG), one of the green tea polyphenols, has chemo preventative properties as studied in various cancer cells [16-18]. Our previous studies have shown that when used in synergistic combination, ascorbic acid, lysine, proline and EGCG exert a significantly pronounced anti-cancer effect than when any of the nutrients are used alone [19].

Conclusion

Despite several treatment options, the patient outcome for mesothelioma is extremely poor, especially once the cancer

has metastasized. Our previous in vitro studies have indicated that NM was able to inhibit growth of MSTO-211H by down-regulating MMP-2 and -9 expression, and up-regulating the MMP inhibitors [4]. The pharmaceutical drugs with pro-apoptotic action act indiscriminately, inducing apoptosis not only in cancer cells but also in healthy cells. These drugs can cause brutal side effects including drug resistance and reduced efficacy to liver failure in patients. In another study, we tested NM to evaluate its effects on normal cells. The normal cells did not show increased apoptosis, proving the cancer cell selective action of NM and thus its being much safer than pharmaceutical drugs [20]. Our other studies have shown as well that MN was ineffective in inducing enzymes in vital organs such as the heart, kidney, and liver, even at high concentrations. This shows that NM is non-toxic [21]. Based on our current study, NM is a safe and effective alternative and should be considered as a supplemental option in the treatment of mesothelioma.

Acknowledgement

The research study was funded by Dr. Rath Foundation (Santa Clara, CA, USA) a non-profit organization. We thank Ms. Catherine Flowers for proofreading assistance.

References

1. American Cancer Society.
2. Antman KH (1981) Clinical presentation and natural history of benign and malignant mesothelioma. *Semin Oncol* 8: 313-320. [Link: https://tinyurl.com/y5wc7brj](https://tinyurl.com/y5wc7brj)
3. Roomi MW, Ivanov V, Kalinovsky T, Niedzwiecki A, Rath M (2006) Inhibition of malignant mesothelioma cell matrix metalloproteinase production and invasion by a novel nutrient mixture. *Exp Lung Res* 32: 69-79. [Link: https://tinyurl.com/y2bsonbe](https://tinyurl.com/y2bsonbe)
4. Roomi MW, Kalinovsky T, Niedzwiecki A, Rath M (2013) Modulation of u-PA, MMPs and their inhibitors by a novel nutrient mixture in human lung cancer and mesothelioma cell lines. *Int J Oncol* 42: 1883-1889. [Link: https://tinyurl.com/y5zfat8s](https://tinyurl.com/y5zfat8s)
5. Basu A, Haldar S (1998) The relationship between Bcl2, Bax and p53: consequences for cell cycle progression and cell death. *Mol Hum Reprod* 4: 1099-1109. [Link: https://tinyurl.com/y2mo9f27](https://tinyurl.com/y2mo9f27)
6. Li J, Yuan J (2008) Caspases in apoptosis and beyond. *Oncogene* 27: 6194-6206. [Link: https://tinyurl.com/y4egjjnz](https://tinyurl.com/y4egjjnz)
7. McIlwain DR, Berger T, Mak TW (2013) Caspase Functions in Cell Death and Disease. *Cold Spring Harb Perspect Biol* 5: a008656. [Link: https://tinyurl.com/y2jbgxex](https://tinyurl.com/y2jbgxex)
8. Bellosillo B, Piqué M, Barragán M, Castaño E, Villamor N, et al. (1998) Aspirin and Salicylate Induce Apoptosis and Activation of Caspases in B-Cell Chronic Lymphocytic Leukemia Cells. *Blood* 92: 1406-1414. [Link: https://tinyurl.com/y6nz9dyb](https://tinyurl.com/y6nz9dyb)
9. Castaño E, Dalmau M, Barragán M, Pueyo G, Bartrons R, et al. (1999) Aspirin induces cell death and caspase-dependent phosphatidylserine externalization in HT-29 human colon adenocarcinoma cells. *Br J Cancer* 81: 294-299. [Link: https://tinyurl.com/y673cwto](https://tinyurl.com/y673cwto)
10. Levine M (1986) New concepts in the biology and biochemistry of ascorbic acid. *N Engl J Med* 314: 892-902. [Link: https://tinyurl.com/y2yaru5cs](https://tinyurl.com/y2yaru5cs)
11. Block G, Menkes M (1989) Ascorbic acid in cancer prevention. In: Moon TE, Micozzi MS. eds. *Diet and cancer prevention. Investigating the role of micronutrients.* New York: Marcel Dekker Inc 3: 41-88. [Link: https://tinyurl.com/y2ydszsf](https://tinyurl.com/y2ydszsf)

12. Block G (1991) Vitamin C and cancer prevention: the epidemiologic evidence. *Am J Clin Nutr* 53 (1 Suppl): 270S-282S. [Link: https://tinyurl.com/y48n88hd](https://tinyurl.com/y48n88hd)
13. Hanck AB (1988) Vitamin C and cancer. *Progress in Clinical and Biological Research* 259: 307-320.
14. Cameron E, Pauling L (1973) Ascorbic acid and the glycosaminoglycans: an orthomolecular approach to cancer and other diseases. *Oncology* 27: 181-192. [Link: https://tinyurl.com/yxd7275f](https://tinyurl.com/yxd7275f)
15. Cameron E, Pauling L (1974) The orthomolecular treatment of cancer I. The role of ascorbic acid in host resistance. *Chem Biol Interact* 9: 273-283. [Link: https://tinyurl.com/y3jnwseq](https://tinyurl.com/y3jnwseq)
16. Valcic S, Timmerman BN, Alberts DS, Wächter GA, Krutzsch M, et al. (1996) Inhibitory effects of six green tea catechins and caffeine on the growth of four selected human tumor cell lines. *Anticancer drugs* 7: 461-468. [Link: https://tinyurl.com/y32fj3u9](https://tinyurl.com/y32fj3u9)
17. Mukhtar H, Ahmed N (2000) Tea polyphenols: Prevention of cancer and optimizing health. *Am J Clin Nutr* 71 (6 Suppl): 1698S-1720S. [Link: https://tinyurl.com/y4kj5fvj](https://tinyurl.com/y4kj5fvj)
18. Yang GY, Liao J, Kim K (1998) Inhibition of growth and induction of apoptosis in human cancer cell lines by tea polyphenols. *Carcinogenesis* 19: 611-616. [Link: https://tinyurl.com/yyl2th56](https://tinyurl.com/yyl2th56)
19. Netke SP, Roomi MW, Ivanov V, Niedzwiecki A, Rath M (2003) A specific combination of ascorbic acid, lysine, proline and epigallocatechin gallate inhibits proliferation of extracellular matrix invasion of various human cancer cell lines. *Research communications in Pharmacology and Toxicology: Emerging Drugs* 2: 37-50.
20. Roomi M, Shanker N, Niedzwiecki A, Rath M (2015) Induction of Apoptosis in the Human Prostate Cancer Cell Line DU-145 by a Novel Micronutrient Formulation. *O J Apo* 4: 11-21. [Link: https://tinyurl.com/yxln6uyw](https://tinyurl.com/yxln6uyw)
21. Roomi MW, Ivanov V, Netke S, Niedzwiecki A, Rath M (2003) Serum markers of the liver, heart, and kidney and lipid profile and histopathology in ODS rats treated with nutrient synergy. *J Am College Nutrit* 22: 477. [Link: https://tinyurl.com/y6orbszv](https://tinyurl.com/y6orbszv)

Discover a bigger Impact and Visibility of your article publication with Peertechz Publications

Highlights

- ❖ Signatory publisher of ORCID
- ❖ Signatory Publisher of DORA (San Francisco Declaration on Research Assessment)
- ❖ Articles archived in worlds' renowned service providers such as Portico, CNKI, AGRIS, TDNet, Base (Bielefeld University Library), CrossRef, Scilit, J-Gate etc.
- ❖ Journals indexed in ICMJE, SHERPA/ROME0, Google Scholar etc.
- ❖ OAI-PMH (Open Archives Initiative Protocol for Metadata Harvesting)
- ❖ Dedicated Editorial Board for every journal
- ❖ Accurate and rapid peer-review process
- ❖ Increased citations of published articles through promotions
- ❖ Reduced timeline for article publication

Submit your articles and experience a new surge in publication services (<https://www.peertechz.com/submission>).

Peertechz journals wishes everlasting success in your every endeavours.

Copyright: © 2019 Roomi MW, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.